



Product Information

Cholesteryl Ester Transfer Protein (CETP), recombinant

Catalog No. R8899

Description	Active, human recombinant CETP, partially purified. Useful as a standardized source of CETP for mass or activity assays, high-throughput screening, or for performing structure-activity relationship (SAR) studies on inhibitor compounds.
Unit Size	~50 µg CETP in 1.2 mg/ml total protein
Source	CHO cells over expressing the full-length human protein.
Storage and Handling	Store at -80° C. Avoid freeze/thaw cycles. Shelf life: 3 years, if stored properly. Wear appropriate gloves, protective clothing and eyewear. Follow safe laboratory practices.

R8899 Cited References

1. Jahangiri A, de Beer MC, Noffsinger V, et al. HDL remodeling during the acute phase response. *Arterioscler Thromb Vasc Biol.* **2009**;29(2):261-267. 10.1161/ATVBAHA.108.178681.
2. Tory, R., Sachs-Barrable, K., Hill, J. S. & Wasan, K. M. Cyclosporine A and Rapamycin induce in vitro cholesteryl ester transfer protein activity, and suppress lipoprotein lipase activity in human plasma. *Int. J. Pharm.* **358**, 219-223 (2008).
3. Deguchi, H., Fernandez, J. A. & Griffin, J. H. Plasma cholesteryl ester transfer protein and blood coagulability. *Thromb. Haemost.* **98**, 1160-1164 (2007).
4. Okamoto, H. *et al.* Cholesteryl ester transfer protein promotes the formation of cholesterol-rich remnant like lipoprotein particles in human plasma. *Clin. Chim. Acta* **375**, 92-98 (2007).
5. Gauthier, A., Lau, P., Zha, X., Milne, R. & McPherson, R. Cholesteryl Ester Transfer Protein Directly Mediates Selective Uptake of High Density Lipoprotein Cholesteryl Esters by the Liver. *Arterioscler. Thromb. Vasc. Biol.* **25**, 2177-2184 (2005).
6. Parathath, S. *et al.* Glycine 420 near the C-terminal transmembrane domain of SR-BI is critical for proper delivery and metabolism of high density lipoprotein cholesteryl ester. *J. Biol. Chem.* **279**, 24976-24985 (2004).
7. Vassiliou, G. & McPherson, R. Role of cholesteryl ester transfer protein in selective uptake of high density lipoprotein cholesteryl esters by adipocytes. *J. Lipid Res.* **45**, 1683-1693 (2004).
8. Connelly, M. A. *et al.* Separation of lipid transport functions by mutations in the extracellular domain of scavenger receptor class B, type I. *J. Biol. Chem.* **278**, 25773-25782 (2003).

Related Products

RB-CETP	Roar CETP Activity Assay Kit
RB-EVAK	Roar Ex Vivo CETP Activity Assay Kit
RB-RPAK	Roar CETP RP Activity Assay Kit

For Research Use Only. Not for Diagnostic or Therapeutic Purposes.

Roar Biomedical, Inc., Audubon Biomedical Center, 3960 Broadway, New York, NY 10032 USA
Tel: +1 (212) 280-2983 ■ Fax: +1 (212) 280-2968 ■ info@roarbiomedical.com ■ www.roarbiomedical.com

©2009 Roar Biomedical, Inc. All rights reserved. This information is subject to change without notice.

Roar rCETP 01W