

R8899 Cited References

Roar Human Recombinant CETP, Active

1. Chen D, Huang X, Zhou H, et al. Discovery of pentacyclic triterpene 3 β -ester derivatives as a new class of cholesterol ester transfer protein inhibitors. *Eur J Med Chem.* **2017**;139:201-213.
2. Kawai Y, et al., inventor "Acid addition salt of substituted pyridine compound." U.S. Patent No. 9,321,747. **2016**.
3. Venancio TM, Machado RM, Castoldi A, et al. CETP lowers TLR4 expression which attenuates the inflammatory response induced by LPS and polymicrobial sepsis. *Mediators Inflamm.* **2016**;2016.
4. Wang J, Bie J, Ghosh S. Intracellular cholesterol transport proteins enhance hydrolysis of HDL-CEs and facilitate elimination of cholesterol into bile. *J Lipid Res.* **2016**;57(9):1712-1719.
5. Xie H, Li Y, Bai C, et al. Discovery of novel N, N-3-phenyl-3-benzylaminopropionanilides as potent inhibitors of cholesteryl ester transfer protein in vivo. *Bioorg Med Chem.* **2016**;24(8):1811-1818.
6. Zhao D, Xie H, Bai C, et al. Design, synthesis and biological evaluation of N, N-3-phenyl-3-benzylaminopropanamide derivatives as novel cholesteryl ester transfer protein inhibitor. *Bioorg Med Chem.* **2016**;24:1589-1597.
7. Pollard RD, Blesso CN, Zabalawi M, et al. Procollagen C-endopeptidase enhancer protein 2 (PCPE2) reduces atherosclerosis in mice by enhancing scavenger receptor class B1 (SR-BI)-mediated high-density lipoprotein (HDL)-cholesteryl ester uptake. *J Biol Chem.* **2015**;290(25):15496-15511.
8. Jahangiri A, de Beer MC, Noffsinger V, et al. HDL remodeling during the acute phase response. *Arterioscler Thromb Vasc Biol.* **2009**;29(2):261-267. doi: 10.1161/ATVBAHA.108.178681.
9. Tory R, Sachs-Barrable K, Hill JS, Wasan KM. Cyclosporine A and rapamycin induce in vitro cholesteryl ester transfer protein activity, and suppress lipoprotein lipase activity in human plasma. *Int J Pharm.* **2008**;358(1-2):219-223.
10. Deguchi H, Fernandez JA, Griffin JH. Plasma cholesteryl ester transfer protein and blood coagulability. *Thromb Haemost.* **2007**;98(6):1160-1164.
11. Okamoto H, Miyai A, Sasase T, et al. Cholesteryl ester transfer protein promotes the formation of cholesterol-rich remnant like lipoprotein particles in human plasma. *Clin Chim Acta.* **2007**;375(1-2):92-98.
12. Gauthier A, Lau P, Zha X, Milne R, McPherson R. Cholesteryl ester transfer protein directly mediates selective uptake of high density lipoprotein cholesteryl esters by the liver. *Arterioscler Thromb Vasc Biol.* **2005**;25(10):2177-2184.
13. Parathath S, Sahoo D, Darlington YF, et al. Glycine 420 near the C-terminal transmembrane domain of SR-BI is critical for proper delivery and metabolism of high density lipoprotein cholesteryl ester. *J Biol Chem.* **2004**;279(24):24976-24985.
14. Vassiliou G, McPherson R. Role of cholesteryl ester transfer protein in selective uptake of high density lipoprotein cholesteryl esters by adipocytes. *J Lipid Res.* **2004**;45(9):1683-1693.
15. Connelly MA, De La Llera-Moya M, Peng Y, Drazul-Schrader D, Rothblat GH, Williams DL. Separation of lipid transport functions by mutations in the extracellular domain of scavenger receptor class B, type I. *J Biol Chem.* **2003**;278(28):25773-25782.